





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 401/12, A61K 31/445		A1	(11) International Publication Number: WO 96/21661
			(43) International Publication Date: 18 July 1996 (18.07.96)
(21) International Application Number: PCT/EP96/00082		(74) Agent: FILLER, W., A.; Glaxo Wellcome plc, Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).	
(22) International Filing Date: 10 January 1996 (10.01.96)			
(30) Priority Data:		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AZ, BY, KZ, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
9500549.2 12 January 1995 (12.01.95) GB 9505639.6 21 March 1995 (21.03.95) GB 9505640.4 21 March 1995 (21.03.95) GB			
(71) Applicant (for all designated States except US): GLAXO GROUP LIMITED [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): ARMOUR, Duncan, Robert [GB/GB]; Glaxo Research and Development Limited, Gunnels Wood Road, Stevenage, Hertsfordshire SG1 2NY (GB). GIBLIN, Gerald, Martin, Paul [GB/GB]; Glaxo Research and Development Limited, Gunnels Wood Road, Stevenage, Hertsfordshire SG1 2NY (GB). PENNELL, Andrew, Michael, Kenneth [GB/GB]; Glaxo Research and Development Limited, Gunnels Wood Road, Stevenage, Hertsfordshire SG1 2NY (GB). SHARRATT, Peter, John [GB/GB]; 46 The Moor, Melbourn, Royston, Hertsfordshire SG8 6ED (GB).			
Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>			

(54) Title: PIPERIDINE DERIVATIVES HAVING TACHYKININ ANTAGONIST ACTIVITY

(57) Abstract

The invention relates to piperidine derivatives of formula (I) wherein R¹ is -O-(CH₂)_pC₃₋₇cycloalkyl, -O-C₁₋₇fluoroalkyl, or -O-(CH₂)_nX; R² is (1); R³ is hydrogen or halogen atom; R⁴ and R⁵ may each independently represent a hydrogen or halogen atom, or a C₁₋₄alkyl, C₁₋₄alkoxy or trifluoromethyl group; X is selected from C(O)-NR⁷R⁸, C(O)R⁹, NR⁷R⁸, SO₂NR⁷R⁸, NHSO₂R⁹, S(O)₂R⁹, OC₁₋₄alkyl, NO₂, CO₂H, CO₂C₁₋₄alkyl, CN or, when n is 2, X may also represent OH, SH or NH₂; R⁶ is a hydrogen atom, a C₁₋₄alkyl, (CH₂)_mcyclopropyl, -S(O)₂C₁₋₄alkyl, phenyl, NR¹⁰R¹¹, CH₂C(O)CF₃, trifluoromethyl, difluoromethyl or cyano group; R⁷ and R⁸ may each independently represent hydrogen atoms or a C₁₋₄alkyl group; R⁹ represents a C₁₋₄alkyl or trifluoromethyl group; R¹⁰ and R¹¹ may each independently represent a hydrogen atom, or a C₂1-4alkyl or acyl group; x represents zero or 1; n is 1 or 2; s represents zero, 1 or 2; m represents zero or 1; p represents zero or 1; and pharmaceutically acceptable salts and solvates thereof, to processes for their preparation, and their use in the treatment of conditions mediated by tachykinins.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Larvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam


PIPERIDINE DERIVATIVES HAVING TACHYKININ ANTAGONIST ACTIVITY

The present invention relates to piperidine derivatives, to processes for their preparation, pharmaceutical compositions containing them and their medical use.

- 5 3-Aminopiperidine derivatives described as having substance P antagonist activity are disclosed in, for example, PCT Patent Applications WO93/00331, WO93/01170 and WO94/13663.


In particular the invention relates to novel compounds which are potent and specific antagonists of tachykinins, including substance P and other neurokinins.

- 10 The present invention provides compounds of formula (I)



wherein R^1 is $-O-(CH_2)_pC_{3-7}$ cycloalkyl, $-O-C_{1-7}$ fluoroalkyl, or $-O-(CH_2)_nX$;

R^2 is



- 15 R^3 is a hydrogen or halogen atom;

R^4 and R^5 may each independently represent a hydrogen or halogen atom, or a C_{1-4} alkyl, C_{1-4} alkoxy or trifluoromethyl group;

X is selected from $C(O)-NR^7R^8$, $C(O)R^9$, NR^7R^8 , $SO_2NR^7R^8$, $NHCO_2R^9$, $S(O)_sR^9$, $OC_{1-4}alkyl$, NO_2 , CO_2H , $CO_2C_{1-4}alkyl$, CN or, when n is 2, X may also represent OH , SH or NH_2 ;

5 R^6 is a hydrogen atom, a $C_{1-4}alkyl$, $(CH_2)_m cyclopropyl$, $-S(O)_sC_{1-4}alkyl$, phenyl, $NR^{10}R^{11}$, $CH_2C(O)CF_3$, trifluoromethyl, difluoromethyl or cyano group;

R^7 and R^8 may each independently represent hydrogen atoms or a $C_{1-4}alkyl$ group;

R^9 represents a $C_{1-4}alkyl$ or trifluoromethyl group;

10 R^{10} and R^{11} may each independently represent a hydrogen atom, or a $C_{1-4}alkyl$ or acyl group;

x represents zero or 1;

n is 1 or 2;

s represents zero, 1 or 2;

m represents zero or 1;

15 p represents zero or 1;

and pharmaceutically acceptable salts and solvates thereof.

20 Suitable pharmaceutically acceptable salts of the compounds of general formula (I) include acid addition salts formed with pharmaceutically acceptable organic or inorganic acids for example, hydrochlorides, hydrobromides, sulphates, alkyl- or arylsulphonates (e.g. methanesulphonates or p-toluenesulphonates), phosphates, acetates, citrates, succinates, tartrates, fumarates and maleates. Dihydrochloride salts are particularly suitable.

25 Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be useful in the preparation of salts useful as intermediates in obtaining the compounds of formula (I) and their pharmaceutically acceptable acid addition salts.


The solvates may, for example, be hydrates.

References hereinafter to a compound according to the invention, includes both compounds of formula (I) and their pharmaceutically acceptable acid addition salts together with pharmaceutically acceptable solvates.

It will be appreciated by those skilled in the art that the compounds of formula (I) contain at least two chiral centres (shown as * in formula (I)) and thus exist in the form of two pairs of optical isomers (i.e. enantiomers) and mixtures thereof including racemic mixtures.

For example the compounds of formula (I) may be either cis isomers, as represented by figures (a) and (b), or trans isomers, as represented by figures (c) and (d), or mixtures thereof.

All of the isomers of the compounds of formula (I) represented by the figures (a) to (d) and mixtures thereof including racemic mixtures are included within the scope of the invention.



The compounds of formula (I) are preferably in the form of their cis isomers (i.e. as represented by figures (a) and (b)). The 2S, 3S isomers (i.e. as represented by figure (b)) are particularly preferred.

Referring to the general formula (I), a C₁₋₄alkoxy group may be a straight chain or branched chain alkoxy group, for example, methoxy, ethoxy, propoxy, prop-2-

oxy, butoxy, but-2-oxy, 2-methylprop-1-oxy or 2-methylprop-2-oxy. A C_{1-4} alkyl group may be a straight chain or branched chain alkyl group and may be, for example, methyl, ethyl, propyl, prop-2-yl, butyl, but-2-yl, 2-methylprop-1-yl or 2-methylprop-2-yl. An $-O-(CH_2)_pC_{3-7}$ cycloalkyl group may be, for example

5 cyclopropyloxy, cyclopropylmethyloxy, cyclobutyloxy, cyclobutylmethyloxy, cyclopentyloxy, cyclohexyloxy or cycloheptyloxy.

Referring to the general formula (I), an $-O-C_{1-7}$ fluoroalkyl group may contain 1, 2 or 3 fluorine atoms and the C_{1-7} alkyl chain may be straight or branched. Examples of suitable groups include fluoromethyloxy, difluoromethyloxy,

10 trifluoromethyloxy and 2,2,2-trifluoroethyloxy.

Referring to the general formula (I), when R^7 , R^8 , or R^9 represents a C_{1-4} alkyl group, this is suitably methyl.

Referring to the general formula (I), a halogen atom may be a fluorine, chlorine, bromine or iodine atom, such as a fluorine, chlorine or bromine atom.


15 Referring to the general formula (I), R^1 is suitably $-O-(CH_2)_nX$ where X is suitably a group selected from $C(O)NHMe$, $C(O)NH_2$, $C(O)NMe_2$, $C(O)Me$, $C(O)CF_3$, $NHMe$, NMe_2 , SO_2NH_2 , SO_2NHMe , SO_2NMe_2 , $NHSO_2Me$, $NHSO_2CF_3$, $S(O)_sMe$ or $S(O)_sCF_3$, for example where s is 2, OMe , NO_2 , CO_2H , CO_2Me , CN or, when n is 2, X is suitably OH , SH or NH_2 .

20 X is preferably OC_{1-4} alkyl, for example OMe .

Referring to the general formula (I), R^1 is suitably a $-O-(CH_2)_p$ -cyclopropyl, $-O-(CH_2)_p$ -cyclopentyl, $-O-C_{1-2}$ fluoroalkyl or $O-(CH_2)_nOC_{1-4}$ alkyl group, such as cyclopropylmethyloxy, cyclopentyloxy, fluoromethyloxy, difluoromethyloxy, trifluoromethyloxy, 2,2,2-trifluoroethyloxy, especially fluoromethyloxy, $O-$

25 $(CH_2)_nOMe$, for example OCH_2OMe or $O-CH_2CH_2OMe$.

Referring to the general formula (I), R^2 is suitably a group



Referring to the general formula (I), when R^6 is an $NR^{10}R^{11}$ group; this is suitably NH_2 , $NH(C_{1-4}alkyl)$, $NHacyl$ i.e. $NHC(O)methyl$, or $N(C_{1-4}alkyl)_2$.

Referring to the general formula (I), when R^6 is a $C_{1-4}alkyl$ group, this is suitably methyl.

- 5 Referring to the general formula (I), when R^2 is a group (A) as defined above, R^6 is suitably a hydrogen atom or a $C_{1-4}alkyl$, e.g. methyl, or a trifluoromethyl group.

R^2 is preferably a group (A) as defined above.


R^3 is preferably a hydrogen atom.

R^4 and R^5 are preferably hydrogen atoms.

- 10 R^6 is preferably a hydrogen atom, a $C_{1-4}alkyl$, e.g. methyl, or a trifluoromethyl group.

x is preferably zero.

A preferred class of compounds of formula (I) are those wherein R^1 is $-O-(CH_2)_pC_{3-7}cycloalkyl$ or $-O-C_{1-7}fluoroalkyl$, as defined hereinbefore, R^2 is



15

where R^6 is a hydrogen atom, a $C_{1-4}alkyl$, e.g. methyl, or trifluoromethyl group, x is zero and R^3 , R^4 and R^5 are each hydrogen atoms.

- 20 A further preferred class of compounds of formula (I) are those wherein R^1 is a cyclopropylmethoxy, cyclopentyloxy, difluoromethyloxy, trifluoromethyloxy, 2,2,2-trifluoroethyloxy or, more preferably, a fluoromethyloxy group, R^2 is a group (A) as defined above, x is zero, R^3 , R^4 and R^5 are hydrogen atoms, and R^6 is a hydrogen atom or a methyl or trifluoromethyl group.

- 25 A preferred class of compounds of formula (I) are those wherein R^1 is a group $-O-(CH_2)_nX$, R^2 is a group (A) as defined above, x is zero and R^6 is a hydrogen atom or a trifluoromethyl, difluoromethyl or cyano group.

An additional preferred class of compounds of formula (I) are those wherein R¹ is -O-(CH₂)_nX where X is a group selected from C(O)NH₂, C(O)NHMe, C(O)NMe₂, C(O)Me, C(O)CF₃, NHMe, NMe₂, SO₂NH₂, SO₂NHMe, SO₂NMe₂, NHSO₂Me, NHSO₂CF₃, S(O)_sMe or S(O)_sCF₃, for example where s is 2, OMe, NO₂, CO₂H, CO₂Me, CN or, when n is 2, X is OH, SH or NH₂, R² is a group (A) as defined above, x is zero, R⁶ is trifluoromethyl, and R³, R⁴ and R⁵ are each hydrogen atoms.

A further preferred class of compounds of formula (I) are those wherein R¹ is O-(CH₂)_nOC₁₋₄alkyl, such as O-(CH₂)_nOMe, for example OCH₂OMe or O-CH₂CH₂OMe, R² is a group (A) as defined above, x is zero, R³, R⁴ and R⁵ are hydrogen atoms, and R⁶ is trifluoromethyl.

Specific compounds according to the invention include:

(2-Cyclopentoxy-5-tetrazol-1-ylbenzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

(2-Cyclopropylmethoxy-5-tetrazol-1-ylbenzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

(2-Cyclopentoxy-5-(5-trifluoromethyltetrazol-1-yl)benzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

(2-Cyclopropylmethoxy-5-(5-trifluoromethyltetrazol-1-yl)benzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

(2-Fluoromethoxy-5-(5-trifluoromethyltetrazol-1-yl)benzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

[2-(2-methoxy-1-ethoxy)-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-([2S,3S]-2-phenylpiperidine-3-yl)amine

and pharmaceutically acceptable salts, especially the dihydrochloride salts, and solvates thereof.

Further specific compounds according to the invention include:

(2-Cyclopentoxy-5-(5-methyltetrazol-1-yl)benzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

(2-Trifluoromethoxy-5-tetrazol-1-ylbenzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

(2-Trifluoromethoxy-5-(5-methyltetrazol-1-yl)benzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

5 (2-Trifluoromethoxy-5-(5-trifluoromethyltetrazol-1-yl)benzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

(2-Difluoromethoxy-5-tetrazol-1-ylbenzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

(2-Difluoromethoxy-5-(5-methyltetrazol-1-yl)benzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

10 (2-Difluoromethoxy-5-(5-trifluoromethyltetrazol-1-yl)benzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

(2-(2,2,2-Trifluoroethoxy)-5-tetrazol-1-ylbenzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

15 (2-(2,2,2-Trifluoroethoxy)-5-(5-methyltetrazol-1-yl)benzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

(2-(2,2,2-Trifluoroethoxy)-5-(5-trifluoromethyltetrazol-1-yl)benzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

(2-Cyclopropylmethoxy-5-(5-methyltetrazol-1-yl)benzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

20 and pharmaceutically acceptable salts and solvates thereof.

The compounds of the invention are antagonists of tachykinins, including substance P and other neurokinins both in vitro and in vivo and are thus of use in the treatment of conditions mediated by tachykinins, including substance P and other neurokinins.

25 The compounds of the invention possess NK₁- receptor binding affinity as determined in vitro by their ability to displace [3H]- substance P (SP) from recombinant human NK₁ receptors expressed in Chinese Hamster Ovary (CHO) cell membranes. CHO membranes (3-5µg protein per tube) were prepared and

incubated with [3H]-SP (0.6 - 0.8nM) at 20°C for 40min. Non-specific binding was defined as that remaining in the presence of 1 μ M (+) CP99,994.

5 The compounds of the invention have been shown to have anti-emetic activity as indicated by for example their ability to inhibit radiation-induced emesis in the ferret. In this model of emesis the onset of retching and vomiting occurs approximately 20 minutes after whole body irradiation (2 Grey \equiv 200 Rads). The test compound is administered (e.g. i.p, p.o., i.v., s.c) immediately after irradiation and its effect on emesis determined by comparison with appropriate controls. Anti-emetic activity may also be demonstrated using other emetogens
10 such as cisplatin and ipecacuanha. Alternatively, the test compounds may be administered before irradiation or before treatment with an emetogen, for example 1.5, 3 or 6 hours before irradiation.

15 The compounds of the invention are potent and specific NK₁ antagonists. Furthermore, they exhibit good oral bioavailability and have an advantageous duration of action.

Compounds of the invention are useful as analgesics in particular they are useful in the treatment of traumatic pain such as postoperative pain; traumatic avulsion pain such as brachial plexus; chronic pain such as arthritic pain such as occurring in osteo-, rheumatoid or psoriatic arthritis; neuropathic pain such as post-herpetic neuralgia, trigeminal neuralgia, segmental or intercostal
20 neuralgia, fibromyalgia, causalgia, peripheral neuropathy, diabetic neuropathy, chemotherapy-induced neuropathy, AIDS related neuropathy, occipital neuralgia, geniculate neuralgia, glossopharyngeal neuralgia, reflex sympathetic dystrophy, phantom limb pain; various forms of headache such as migraine, acute or chronic tension headache, temporomandibular pain, maxillary sinus pain, cluster headache; odontalgia; cancer pain; pain of visceral origin; gastrointestinal pain; nerve entrapment pain; sport's injury pain; dysmennorrhoea; menstrual pain; meningitis; arachnoiditis; musculoskeletal pain; low back pain e.g. spinal stenosis; prolapsed disc; sciatica; angina;
25 ankylosing spondylitis; gout; burns; scar pain; itch; and thalamic pain such as post stroke thalamic pain.
30

Compounds of the invention are also useful as antiinflammatory agents in particular they are useful in the treatment of inflammation in asthma, influenza, chronic bronchitis and rheumatoid arthritis; in the treatment of inflammatory

diseases of the gastrointestinal tract such as Crohn's disease, ulcerative colitis, inflammatory bowel disease, non-steroidal anti-inflammatory drug induced damage and inflammatory and secretory effects of bacterial infection, e.g. due to *Clostridium difficile*; inflammatory diseases of the skin such as herpes and eczema; inflammatory diseases of the bladder such as cystitis and urge (i.e. urinary) incontinence; and eye and dental inflammation, e.g. gingivitis and periodontitis.

Compounds of the invention are also useful in the treatment of allergic disorders in particular allergic disorders of the skin such as urticaria, and allergic disorders of the airways such as rhinitis.

Compounds of the invention may also be useful in the treatment of CNS disorders in particular psychoses such as schizophrenia, mania or dementia; cognitive disorders e.g. Alzheimer's disease; anxiety; AIDS related dementia; diabetic neuropathy; multiple sclerosis; depression; Parkinson's disease; and dependency on drugs or substances of abuse; and also the compounds of the invention may act as myorelaxants and antispasmodics.

Compounds of the invention are also useful in the treatment of emesis, i.e. nausea, retching and vomiting. Emesis includes acute emesis, delayed or late emesis and anticipatory emesis. The compounds of the invention, are useful in the treatment of emesis however induced. For example, emesis may be induced by drugs such as cancer chemotherapeutic agents such as alkylating agents, e.g. cyclophosphamide, carmustine, lomustine and chlorambucil; cytotoxic antibiotics, e.g. dactinomycin, doxorubicin, mitomycin-C and bleomycin; anti-metabolites, e.g. cytarabine, methotrexate and 5-fluorouracil; vinca alkaloids, e.g. etoposide, vinblastine and vincristine; and others such as cisplatin, dacarbazine, procarbazine and hydroxyurea; and combinations thereof; radiation sickness; radiation therapy, e.g. irradiation of the thorax or abdomen, such as in the treatment of cancer; carcinoid syndrome; poisons; toxins such as toxins caused by metabolic disorders or by infection, e.g. gastritis, or released during bacterial or viral gastrointestinal infection; pregnancy; vestibular disorders, such as motion sickness, vertigo, dizziness and Meniere's disease; post-operative sickness; dialysis-induced emesis; prostaglandin-induced emesis; gastrointestinal obstruction; reduced gastrointestinal motility; visceral pain, e.g. myocardial infarction or peritonitis; migraine; increased intracranial

pressure; decreased intracranial pressure (e.g. altitude sickness); opioid analgesics, such as morphine; gastro-oesophageal reflux disease, acid indigestion, over-indulgence of food or drink, acid stomach, sour stomach, waterbrash/regurgitation, heartburn, such as episodic heartburn, nocturnal heartburn, and meal-induced heartburn and dyspepsia; organ failure, such as in terminal illness; AIDS and AIDS-related conditions, and treatments thereof; and cyclic vomiting syndrome.

Compounds of the invention are also useful in the treatment of gastrointestinal disorders such as irritable bowel syndrome; skin disorders such as psoriasis, pruritis and sunburn; vasospastic diseases such as angina, vascular headache and Reynaud's disease; cerebral ischaemia such as cerebral vasospasm following subarachnoid haemorrhage; fibrosing and collagen diseases such as scleroderma and eosinophilic fasciitis; disorders related to immune enhancement or suppression such as systemic lupus erythematosus and rheumatic diseases such as fibrositis; and cough.

The invention therefore provides a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof for use in therapy, in particular in human medicine.

There is also provided as a further aspect of the invention the use of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof in the preparation of a medicament for use in the treatment of conditions mediated by tachykinins, including substance P and other neurokinins.

In an alternative or further aspect there is provided a method for the treatment of a mammal, including man, in particular in the treatment of conditions mediated by tachykinins, including substance P and other neurokinins, comprising administration of an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

It will be appreciated that reference to treatment is intended to include prophylaxis as well as the alleviation of established symptoms. Compounds of formula (I) may be administered as the raw chemical but the active ingredient is preferably presented as a pharmaceutical formulation.

Accordingly, the invention also provides a pharmaceutical composition which comprises at least one compound of formula (I) or a pharmaceutically acceptable salt thereof and formulated for administration by any convenient route. Such compositions are preferably in a form adapted for use in medicine, in particular human medicine, and can conveniently be formulated in a conventional manner using one or more pharmaceutically acceptable carriers or excipients.

Thus compounds of formula (I) may be formulated for oral, buccal, parenteral, topical (including ophthalmic and nasal), transdermal, depot or rectal administration or in a form suitable for administration by inhalation or insufflation (either through the mouth or nose).

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g. magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium starch glycollate); or wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g. sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g. lecithin or acacia); non-aqueous vehicles (e.g. almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g. methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, flavouring, colouring and sweetening agents as appropriate.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration the composition may take the form of tablets or lozenges formulated in conventional manner.

5 The compounds of the invention may be formulated for parenteral administration by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form e.g. in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

10 The compounds of the invention may be formulated for topical administration in the form of ointments, creams, gels, lotions, pessaries, aerosols or drops (e.g. eye, ear or nose drops). Ointments and creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Ointments for administration to the eye may be manufactured in a sterile manner using sterilised components.

15 Lotions may be formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilising agents, dispersing agents, suspending agents, thickening agents, or colouring agents. Drops may be formulated with an aqueous or non aqueous base also comprising one or more dispersing agents, stabilising agents, solubilising agents or suspending agents.
20 They may also contain a preservative.

For transdermal administration the compounds according to the invention may be formulated as creams, gels, ointments or lotions or as a transdermal patch. Such compositions may for example be formulated with an aqueous or oily base with the addition of suitable thickening, gelling, emulsifying, stabilising, dispersing, suspending, and/or colouring agents.
25

The compounds of the invention may also be formulated in rectal compositions such as suppositories or retention enemas, e.g. containing conventional suppository bases such as cocoa butter or other glycerides.

30 The compounds of the invention may also be formulated as depot preparations. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds of the invention may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an

acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.


For intranasal administration, the compounds of the invention may be formulated as solutions for administration via a suitable metered or unit dose device or alternatively as a powder mix with a suitable carrier for administration using a suitable delivery device.

The compositions may contain from 0.1% upwards, e.g. 0.1 - 99% of the active material, depending on the method of administration. A proposed dose of the compounds of the invention is 0.05 mg/kg to about 400 mg/kg bodyweight per day e.g. 0.05mg/kg to 5mg/kg per day. It will be appreciated that it may be necessary to make routine variations to the dosage, depending on the age and condition of the patient and the precise dosage will be ultimately at the discretion of the attendant physician or veterinarian. The dosage will also depend on the route of administration and the particular compound selected.


The compounds of formula (I) may, if desired, be administered with one or more therapeutic agents and formulated for administration by any convenient route in a conventional manner. Appropriate doses will be readily appreciated by those skilled in the art. For example, the compounds of formula (I) may be administered in combination with a systematic anti-inflammatory corticosteroid such as methyl prednisolone or dexamethasone, or a 5HT₃ antagonist such as ondansetron, granisetron or metoclopramide. The compounds of formula (I) may also be administered in combination with sympathomimetics, such as ephedrine, pseudoephedrine and oxymetazoline, or the compounds may be administered with conventional analgesics, such as non-steroidal antiinflammatory drugs (NSAIDs), opioids or local anaesthetics. Compounds which are specific antagonists at NK₁ receptors, such as the compounds of formula (I), may be administered in combination with compounds which are specific antagonists at NK₂ receptors.

Compounds of formula (I), and salts and solvates thereof, may be prepared by the general methods outlined hereinafter. In the following description, the groups R¹, R², R³, R⁴ and R⁵ and x are as previously defined for compounds of formula (I) unless otherwise stated.

According to a first general process (A), a compound of formula (I) may be prepared by reacting a compound of formula (II):



with a compound of formula (III)



to form the intermediate imine, which may be isolated if required,

followed by reduction of the imine using a suitable reducing agent such as a metal hydride, for example a borane hydride, alane hydride or a metal hydride complex like lithium aluminum hydride or sodium borohydride, or an organo-metallic complex such as borane- methyl sulphide, 9-borabicyclononane (9-BBN), triethylsilane, sodium triacetoxyborohydride, sodium cyanoborohydride and the like. Alternatively, catalytic hydrogenation may be used, for example using a platinum catalyst in a suitable solvent e.g. ethanol.


The condensation reaction conveniently takes place in a suitable solvent such as an alcohol (e.g. methanol), an aromatic hydrocarbon (e.g. benzene, toluene or xylene) or a chlorinated hydrocarbon (e.g. dichloromethane or dichloroethane) at a temperature ranging from ambient to the reflux temperature of the reaction mixture. The reaction preferably takes place in the presence of a catalytic amount of a suitable acidic condensing agent such as p-toluenesulphonic acid or acetic acid and/or a dehydrating agent such as molecular sieves, or the reaction may take place under Dean-Stark conditions.

The reduction step conveniently takes place in a suitable solvent such as acetonitrile, dimethylformamide, benzene, chlorinated hydrocarbons such as dichloromethane or dichloroethane, ethers such as diethyl ether, tetrahydrofuran, dioxane and 1,2-dimethoxyethane and alcohols such as ethanol at a temperature ranging from 0°C to the reflux temperature of the reaction mixture.

Process (A) may also take place in one step without isolation of the intermediate imine if the condensation reaction takes place in the presence of sodium cyanoborohydride or sodium triacetoxyborohydride. Further reduction is therefore unnecessary in this case.


When carrying out process (A) where R^2 is a group (C) as defined hereinbefore, R^6 is preferably a C_{1-4} alkyl group.

Compounds of formula (II) may be prepared by reducing compounds of formula (IV)



under suitable reducing conditions, such as catalytic hydrogenation, for example using a platinum catalyst, e.g. platinum (IV) oxide, in a suitable solvent like ethanol, preferably in the presence of concentrated hydrochloric acid.

Compounds of formula (IV) may be prepared by reacting 2-chloro-3-nitropyridine with a compound of formula (V)




in the presence of a palladium (0) catalyst such as tetrakis (triphenyl phosphine) palladium (0). The reaction suitably takes place in the presence of

a solvent such as an ether, e.g. dimethoxyethane, at an elevated temperature and preferably in the presence of a base such as sodium carbonate.


Compounds of formula (V) may be prepared by reacting the corresponding bromo-compounds under Grignard conditions followed by reaction with triisopropylborate.

Alternatively, compounds of formula (II) may be prepared by reducing compounds of formula (VI)




under suitable reducing conditions, for example using a metal hydride complex such as sodium borohydride in the presence of zirconium (IV) chloride in a suitable solvent such as tetrahydrofuran.

Compounds of formula (VI) may be prepared by reacting compounds of formula (VII)




with hydroxylamine hydrochloride in the presence of pyridine.

Compounds of formula (VII) may be prepared by reacting compounds of formula (VIII)




with ozone in the presence of potassium t-butoxide in a suitable solvent such as a mixture of dichloromethane and methanol.

Compounds of formula (VIII) may be prepared by reacting compounds of formula (IX)



with methyl-4-nitrobutyrate and ammonium acetate, in a suitable solvent such as an alcohol, e.g. ethanol at elevated temperature.

Compounds of formula (III), except where R^1 represents cyclopropyloxy, cyclobutyloxy or trifluoromethyloxy, may be prepared by reacting compounds of formula (X)




with an appropriate alkylating agent, such as $(CH_2)_pC_{3-7}$ -cycloalkyl bromide or iodide or C_{1-7} -fluoroalkyl chloride, bromide or iodide, in the presence of a base such as potassium carbonate.

Compounds of formula (III) where R^1 represents 2,2,2-trifluoroethyloxy may be prepared by reacting compounds of formula (X) with $CF_3CH_2OSO_2R^*$, where R^* represents methyl, paratolyl or trifluoromethyl, in the presence of hexamethylphosphoramide and potassium carbonate at elevated temperature.

Compounds of formula (III) where R^1 represents trifluoromethyloxy may be prepared by reacting compounds of formula (X), or a protected aldehyde thereof, with carbondisulphide and methyl iodide to give the xanthate derivative. The xanthate derivative is then treated with hydrogenfluoride-pyridine complex and deprotected if necessary, to give the required compound of formula (III).

Compounds of formula (III) where R^1 represents cyclopropyloxy may be prepared by reacting compounds of formula (X), or a protected aldehyde thereof, with ethylvinyl ether in the presence of mercury acetate. The resulting vinyl derivative is treated with diiodomethane and diethylzinc and deprotected if necessary, to give the required compound of formula (III).


Compounds of formula (III) where R^1 represents cyclobutyloxy may be prepared by reacting compounds of formula (XI)



where R^b is a suitable leaving group, such as CF_3SO_3 , or a protected aldehyde thereof, with cyclobutanol in the presence of a base, such as sodium hydride, and a solvent such as dimethylformamide, followed by deprotection where necessary.

Compounds of formula (XI) may be prepared by conventional methods. For example, compounds of formula (XI) where R^b is CF_3SO_3 may be prepared by reacting compounds of formula (X) with trifluoromethanesulphonic anhydride in the presence of a base, such as triethylamine.

Compounds of formula (X) may be prepared by reacting compounds of formula (XII)




with hexamethylenetetramine in the presence of trifluoroacetic acid at elevated temperature.

Compounds of formula (XII) where R^2 is a group (A) as defined hereinbefore and x is zero may be prepared by reacting the appropriate p-hydroxyaniline, or a protected derivative thereof, with compounds of formula (XIII)



- 5 (where R^9 is methyl or ethyl), for example triethylorthoacetate, in acetic acid followed by reaction with sodium azide at elevated temperature and deprotection if necessary.


Compounds of formula (XII) where R^2 is a group (A) as defined hereinbefore and x is zero may also be prepared by reacting a compound of formula (XIV)



10

or a protected derivative thereof, with sodium azide in acetic acid at elevated temperature, followed by deprotection if necessary.

Compounds of formula (XIV) may be prepared by reacting a compound of formula (XV)



15


or a protected derivative thereof, with resin-supported triphenylphosphine in carbon tetrachloride at elevated temperature.

- 20 Compounds of formula (XV) may be prepared by reacting the appropriate p-hydroxyaniline, or a protected derivative thereof, with the appropriate acid chloride or anhydride, i.e. R^6-COCl or $R^6-CO.O.CO-R^6$, for example trifluoroacetic anhydride, cyclopropane carbonyl chloride, or difluoroacetic

anhydride (or the activated acid may be prepared in situ from difluoroacetic acid and e.g. dicyclohexylcarbodiimide).

Compounds of formula (XII) where R^2 is a group (A) as defined hereinbefore and x is zero, or protected derivatives thereof, may alternatively be prepared by reacting compounds of formula (XV), or protected derivatives thereof, with an acid anhydride such as trifluoroacetic anhydride or trifluoromethane sulfonic anhydride and sodium azide in acetonitrile.

Compounds of formula (III) where R^2 is a group (A) as defined hereinbefore, x is zero and R^6 is $-NH_2$ may alternatively be prepared by reacting compounds of formula (XVI)




or a protected derivative thereof, with ammonium chloride and sodium azide at elevated temperature, suitably in a solvent such as dimethylformamide, followed by deprotection where required.

Compounds of formula (XVI) may be prepared by reacting compounds of formula (III) where R^2 is a group (A) as defined hereinbefore, x is zero and R^6 is hydrogen, or a protected derivative thereof, with n-butyl lithium in a suitable solvent such as tetrahydrofuran.

Compounds of formula (III), or protected derivatives thereof, where R^6 represents one group may be converted into other compounds of formula (III), or protected derivatives thereof, where R^6 represents a different group using conventional procedures, such as alkylation, acylation or oxidation.


Compounds of formula (III) may alternatively be prepared by oxidising compounds of formula (XVII)

21



with a suitable oxidising agent such as manganese dioxide in a suitable solvent such as an ether, e.g. tetrahydrofuran, at elevated temperature.

- 5 Compounds of formula (XVII), except where R⁶ is a cyano group, may be prepared by reducing compounds of formula (XVIII)




with a suitable reducing agent such as a metal hydride complex such as lithium borohydride in a suitable solvent such as an ether, e.g. tetrahydrofuran, or an alcohol, e.g. ethanol, or a mixture thereof.

- 10 Compounds of formula (XVIII) where R² is a group (A) as defined hereinbefore and x is zero may be prepared from the corresponding 2-alkoxy-5-amino benzoic acid methyl ester by reacting with compounds of formula (XIII) as defined above, e.g. triethyl orthoformate, and sodium azide in glacial acetic acid and dimethylform- amide at elevated temperature.

- 15 Suitable 2-alkoxy-5-amino benzoic acid methyl esters are either known or may be prepared according to methods known for the preparation of known compounds e.g. the method described by Bergman *et al* in Can.J.Chem., (1973), 51, 162-70.


- 20 Compounds of formula (III) where R² is a group (A) or (B) as defined hereinbefore and x is 1, may be prepared by reacting compounds of formula (XIX)

22



(where hal is a halogen, e.g. bromine or chlorine, atom) with tetrazole in the presence of a base such as triethylamine or potassium carbonate in a suitable solvent such as dichloromethane or dimethylformamide.

- 5 Compounds of formula (XIX) where x is 1 may be prepared by reacting compounds of formula (XX)




- 10 or a protected derivative thereof, with a carbon tetrahalide, e.g. carbon tetrabromide, in the presence of triphenylphosphine and a suitable solvent such as ether, followed by deprotection where required.


Compounds of formula (XX) may be prepared by reduction of the corresponding aldehydes after protection of the aldehyde group ortho to R¹.

- 15 Compounds of formula (XII) where R² is a group (B) and x is zero may be prepared by reacting the appropriate 1-fluoro-4-nitrobenzene with 1H-tetrazole in a suitable solvent at elevated temperature, followed by reduction of the nitro group by catalytic hydrogenation, followed by conversion of the resulting amino function into an alcohol function using nitrous acid.

Compounds of formula (III) where R² is a group (C) as defined hereinbefore may be prepared by reacting a compound of formula (XXI)




with a compound of formula (XXII)



5 in the presence of a palladium (O) catalyst such as tetrakis (triphenylphosphine) palladium (O) in a suitable solvent such as an ether (e.g. dimethoxyethane) at an elevated temperature.

Compounds of formula (XXI) may be prepared according to similar methods for the preparation of compounds of formula (V) above.

10 Alternatively, compounds of formula (III) where R² is a group (C) as defined hereinbefore, x is zero and R⁶ is hydrogen may be prepared by reacting compounds of formula (XXIII)




or a protected derivative thereof, with tributyltinazide at elevated temperature, followed by deprotection where necessary.

15 Compounds of formula (XXIII) may be prepared from the appropriate p-hydroxybenzonitrile and hexamethylenetetramine as described above for the preparation of compounds of formula (X) from compounds (XII).

According to a further general process (B), compounds of formula (I) where R² is a group (A) as defined hereinbefore, x is zero and R⁶ is -NH₂ may be prepared

20 by reacting compounds of formula (XXIV)


24



with ammonium chloride and sodium azide under conditions as described above for the preparation of compounds of formula (III) from compounds of formula (XVI).

- 5 Compounds of formula (XXIV) may be prepared by reacting compounds of formula (XVI) with compounds of formula (III) under conditions as described above for process (A).

According to a further general process (C), compounds of formula (I) may be prepared by reduction of compounds of formula (XXV)




10

with a suitable reducing agent, such as a metal hydride, for example a borane hydride, in a suitable solvent such as an ether, e.g. tetrahydrofuran, at ambient temperature.

- 15 Compounds of formula (XXV) may be prepared by reacting compounds of formula (III) with compounds of formula (XXVI)


25



under conditions as described above for process (A).

Compounds of formula (XXVI) are either known or may be prepared according to methods known for the preparation of known compounds, for example according to the method described in European Patent Application No. EP-A-0436334, incorporated herein by reference.


According to a further general process (D) compounds of formula (I), excluding compounds where R^1 represents cyclopropyloxy, cyclobutyloxy and trifluoromethyloxy, may be prepared by reacting a compound of formula (XXVII)



or an amino-protected derivative thereof, with $L-(CH_2)_pC_{3-7}\text{cycloalkyl}$ or $L-C_{1-7}\text{fluoroalkyl}$ where p is as defined hereinbefore and L represents a suitable leaving group such as iodine, bromine, chlorine, OSO_2R^c , where R^c represents methyl, paratolyl or trifluoromethyl, in the presence of a base, such as potassium carbonate, and a suitable solvent, such as dimethylformamide, at a temperature between 0 and 100°C, e.g. at room temperature, followed by deprotection where necessary.

Compounds of formula (XXVII) may be prepared by reacting compounds of formula (XXVIII)

26




where R^{10} is C_{1-4} alkyl, e.g. methyl, with borane-methyl sulphide complex in a suitable solvent, such as dichloromethane, at elevated temperature.

5

Compounds of formula (XXVIII) may be prepared according to similar methods described herein for the preparation of compounds of formula (I) and those methods described in PCT patent application no. PCT/EP94/03129.

According to a further general process (E) compounds of formula (I) where R^2 is a group A as defined above and R^6 represents cyano may be prepared by reacting a compound of formula (XXIX)



10

where L represents a suitable leaving group, such as a methylsulphonyl group, with potassium cyanide. The reaction conveniently takes place at elevated temperature and in the presence of a suitable solvent, such as an alcohol, e.g. ethanol.

15

Compounds of formula (XXIX) may be prepared by the methods described in processes (A) to (D) described hereinbefore from the appropriately substituted

intermediates and also from those methods described in PCT patent application no. PCT/EP94/03129.

5 Suitable protecting groups for the hydroxyl function include benzyl groups which may be introduced and removed according to conventional procedures. For example deprotection may be effected by catalytic hydrogenation.

Aldehyde functions may be protected as acetals which may be introduced and removed according to conventional procedures. For example, deprotection may be effected by acid hydrolysis.

10 Amino functions may be protected by, for example, a t-butyl carbamate (BOC) group or a benzyl group which may be introduced and removed according to conventional procedures. For example, deprotection of BOC groups may be effected by acid hydrolysis.

Compounds of formulae (III), (XVI), (XVII), (XVIII), (XXIV), (XXV), (XXVII) and (XXIX) are novel and therefore form a further feature of the invention.

15 Where it is desired to isolate a compound of formula (I) as a salt, for example a pharmaceutically acceptable salt, this may be achieved by reacting the compound of formula (I) in the form of the free base with an appropriate amount of suitable acid and in a suitable solvent such as an alcohol (e.g. ethanol or methanol), an ester (e.g. ethyl acetate) or an ether (e.g. diethyl ether or
20 tetrahydrofuran).

Pharmaceutically acceptable salts may also be prepared from other salts, including other pharmaceutically acceptable salts, of the compound of formula (I) using conventional methods.

25 The compounds of formula (I) may readily be isolated in association with solvent molecules by crystallisation from or evaporation of an appropriate solvent to give the corresponding solvates.

When a specific enantiomer of a compound of general formula (I) is required, this may be obtained for example by resolution of a corresponding enantiomeric mixture of a compound of formula (I) using conventional methods.

Thus, in one example an appropriate optically active acid may be used to form salts with the enantiomeric mixture of a compound of general formula (I). The resulting mixture of isomeric salts may be separated for example by fractional crystallisation, into the diastereoisomeric salts from which the required enantiomer of a compound of general formula (I) may be isolated by conversion into the required free base.

Alternatively, enantiomers of a compound of general formula (I) may be synthesised from the appropriate optically active intermediates using any of the general processes described herein.

A particularly suitable route for the preparation of optically active intermediates of formula (II) from the enantiomeric mixture thereof is by fractional crystallisation using (2R, 3R)-bis-(4-methyl-benzoyloxy)-succinic acid. Thus, the cis (S,S) form of intermediate (II) may be obtained from an enantiomeric mixture thereof (e.g. the racemic mixture) by fractional crystallisation with (2R, 3R)-bis-(4-methyl-benzoyloxy)-succinic acid in a suitable solvent, such as an aqueous alcohol, e.g. aqueous ethanol, isolating the resulting salt and converting it into the corresponding optically active free base by conventional procedures for example using aqueous ammonia.

Alternative methods for preparing and resolving 2-phenyl-3-aminopiperidine are described in PCT patent application no. WO 94/27966, incorporated herein by reference.

Specific enantiomers of a compound of formula (I) may also be obtained by chromatography of the corresponding enantiomeric mixture on a chiral column, for example by chiral preparative h.p.l.c.

Specific diastereoisomers of a compound of general formula (I) may be obtained by conventional methods for example, by synthesis from an appropriate asymmetric starting material using any of the processes described herein, or by conversion of a mixture of isomers of a compound of general formula (I) into appropriate diastereoisomeric derivatives e.g. salts which can then be separated by conventional means e.g. by chromatography or by fractional crystallisation. Alternatively, the diastereoisomers may be separable without the need for further derivatization.

Standard resolving methods are described for example in 'Stereochemistry of Carbon Compounds' by E.L. Eliel (McGraw Hill, 1962) and 'Tables of Resolving Agents' by S.H. Wilen.

5 The various general methods described above may be useful for the introduction of the desired groups at any stage in the stepwise formation of the required compound, and it will be appreciated that these general methods can be combined in different ways in such multi-stage processes. The sequence of the reactions in multi-stage processes should of course be chosen so that the reaction conditions used do not affect groups in the molecule which are desired
10 in the final product.

The invention is further illustrated by the following Intermediates and Examples which are not intended as a limitation of the invention. All temperatures are in °C. Flash column chromatography (FCC) was carried out on silica (Merck 9385). The following abbreviations are used: ether-diethyl ether.

15 Intermediate 1

4-Tetrazol-1-yl-phenol

To a stirred solution of p-amino-phenol (0.1mol) in glacial acetic acid (140ml) at 70-75° under nitrogen atmosphere was added triethylorthoformate (0.1mol).
20 The mixture was stirred at this temperature for 4h, then, sodium azide (0.32mol) was added portionwise and the reaction was continued for 18h, cooled to room temperature and poured into ice water (400ml) and extracted with diethyl ether (3x400ml) and ethyl acetate (1x400ml), dried (MgSO₄), filtered and concentrated to give a dark brown residue which was triturated with 200ml of a mixture of
25 ethanol:diethyl ether (1:1 v/v) and filtered to afford the title compound in 30% yield.

T.l.c. (ether) R_f 0.65

Intermediate 2

2-Hydroxy-5-tetrazol-1-yl-benzaldehyde

30 A solution of 4-tetrazol-1-yl-phenol (0.01mol) in trifluoroacetic acid (20ml) and hexamethylenetetramine (0.04mol) was heated at 70° for 18h, cooled to room

temperature and quenched with 2N solution of sulfuric acid (50ml). The mixture was extracted with ethyl acetate (3x100ml), dried (MgSO_4), filtered and concentrated to give a residue which was purified by FCC (dichloromethane/methanol (9:1)) to afford the title compound in 30% yield.

5 T.l.c. (dichloromethane/methanol (9:1)) Rf 0.6

Intermediate 3

2-Cyclopentoxo-5-tetrazol-1-ylbenzaldehyde

10 A mixture of 2-hydroxy-5-tetrazol-1-yl-benzaldehyde (500mg) and potassium carbonate (431mg) in dimethylformamide (5ml) at 20° was treated with cyclopentyl bromide (0.33ml). After 24h water (20ml) was added and the mixture extracted with ethyl acetate (2 x 25ml). The organics were washed with water (25ml) and brine (25ml) and dried (Na_2SO_4). Evaporation gave a yellow solid which was purified by FCC (ethyl acetate/cyclohexane (2:1)) to give the title compound, as a yellow solid (334mg).

15 $\nu_{\text{Max}}(\text{KBr}) 1677\text{cm}^{-1}$

Similarly prepared:-

Intermediate 4

2-Cyclopropylmethoxy-5-tetrazol-1-ylbenzaldehyde

20 From cyclopropyl methyl bromide (0.3ml) to give the title compound, as a white solid (164mg).

$\nu_{\text{max}}(\text{KBr}) 1681\text{cm}^{-1}$

Intermediate 5

N-(4-Benzyloxy-phenyl)-2,2,2-trifluoro-acetamide

25 A mixture of 4-benzyloxyaniline hydrochloride (0.19mol) in dichloromethane (750ml) at 0° under nitrogen was treated dropwise with trifluoroacetic anhydride (27.6ml) then triethylamine (60ml). After 24h the mixture was poured into t-butyl methyl ether (1.5l) and was washed with 2N hydrochloric acid (1l). The organic phase was dried (MgSO_4) and evaporated in vacuo to give the title compound as a white solid (52.3g).

30 T.l.c. (cyclohexane/ethylacetate (9:1)) Rf 0.36.

Intermediate 6(4-Benzyloxy-phenyl)-(1-chloro-2,2,2-trifluoro-ethylidene)-amine

A mixture of resin-supported triphenylphosphine (3mmol triphenylphosphine/g resin; 58.6g) and N-(4-benzyloxy-phenyl)-2,2,2-trifluoro-acetamide (20.8g) in carbon tetrachloride (800ml) was heated to reflux under nitrogen for 18h. The mixture was allowed to cool then filtered, washing the resin with dichloromethane (1l) and ether (1l). The organics were concentrated in vacuo to give the title compound as a yellow solid (20.7g).

T.l.c. (Cyclohexane/ethyl acetate (9:1)) Rf 0.81

Intermediate 71-(4-Benzyloxy-phenyl)-5-trifluoromethyl-1H-tetrazole

(4-Benzyloxy-phenyl)-(1-chloro-2,2,2-trifluoro-ethylidene)-amine (66mmol) was added to a stirred flask of glacial acetic acid (250ml) at 70° under nitrogen. After 4 min sodium azide (210mmol) was added and heating was continued for 3h. After cooling the mixture was filtered, the filtrate poured into water (750ml) then extracted with dichloromethane (500ml x3). The combined organic extracts were dried (Na₂SO₄) and evaporated in vacuo. Purification by FCC using hexane-ethyl acetate (19:1) gave the title compound as a white solid (14.5g).

T.l.c. (Cyclohexane/ethyl acetate (19:1)) Rf 0.22

Intermediate 84-(5-Trifluoromethyl-tetrazol-1-yl)-phenol

A solution of 1-(4-benzyloxy-phenyl)-5-trifluoromethyl-1H-tetrazole (45.3mmol) in ethanol (100ml) and tetrahydrofuran (100ml) was hydrogenated at room temperature and atmospheric pressure over 10% palladium-carbon catalyst (6g). After 2h, the mixture was filtered and the filtrate was evaporated to give the title compound (10.4g) as a cream solid.

T.l.c. (Dichloromethane/ethanol/ammonia(200:8:1)) Rf 0.3.

Intermediate 92-Hydroxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzaldehyde

A solution of 4-(5-trifluoromethyl-tetrazol-1-yl)-phenol (45mmol) in trifluoroacetic acid (200ml) and hexamethylenetetramine (186.9mmol) was heated at 100°

under nitrogen for 16h. The reaction mixture was quenched with 2N sulphuric acid (250ml) and extracted with ether (3x250ml). The combined organics were dried (Na_2SO_4) and evaporated to give a dark yellow oil. Purification by FCC (hexane/ether (2:1)) afforded the title compound (8.8g) as a pale yellow solid.

5 T.l.c. (hexane/ether (2:1)) Rf 0.36

Intermediate 10

2-Methoxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzaldehyde

10 A mixture of 2-hydroxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzaldehyde (1.56mmol), potassium carbonate (7.8mmol) and methyl iodide (7.8mmol), in acetone (25ml) was stirred for 18h at 23° under nitrogen. Water (150ml) was added and the mixture extracted with diethyl ether (3x50ml). The combined organic extracts were washed with saturated brine, dried (Na_2SO_4) and evaporated to give the title compound as a yellow solid (0.48g).

T.l.c. (ether/hexane (2:1)) Rf 0.38.

15

Intermediate 11

[2-Methoxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-([2S,3S]-2-phenyl-piperidin-3-yl)-amine dihydrochloride

20 A mixture of [2S]-phenyl-piperidin-[3S]-ylamine (1.14mmol), 2-methoxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzaldehyde (1.2mmol), sodium triacetoxyborohydride (2.37mmol) and acetic acid (3 drops) in dichloromethane (25ml) was stirred at 23° under nitrogen for 64h. 2N sodium carbonate solution (50ml) was added and the mixture extracted with dichloromethane (3x25ml). The combined extracts were washed with saturated brine (50ml), dried (MgSO_4)
25 and evaporated. Purification by FCC with dichloromethane/ethanol/ammonia (400:10:1 → 100:10:1) gave a colourless viscous oil. This was dissolved in methanol (10ml) and treated with 2N ethereal hydrogen chloride (~10ml). Evaporation in vacuo and trituration with i-propyl acetate gave the title compound as a white solid (210mg).

30 T.l.c. (Dichloromethane/ethanol/ammonia (200:10:1)) Rf 0.39.

Optical Rotation (c 3mg/ml. water) +50.35°.

Intermediate 12[2-Hydroxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-([2S,3S]-2-phenylpiperidin-3-yl)amine

5 A suspension of [2-methoxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-([2S,3S]-2-phenylpiperidin-3-yl)amine dihydrochloride (1.01g, 2mmol) in chloroform (20ml) was stirred at room temperature and treated with triethylamine, (0.58ml, 406mg, 4mmol) to give a clear solution. This was then treated slowly with a solution of boron tribromide: dimethylsulfide complex (20ml of 1M solution in dichloromethane; 20mmol) and the reaction mixture was refluxed under nitrogen
10 for 3 days. Further reagent (10ml, 10mmol) was then added and after a further 3 days the mixture was cooled in an ice-bath, stirred and treated with methanol (40ml). The mixture was evaporated to dryness. This treatment was repeated twice further. The resulting oil was partitioned between ethyl acetate (100ml) and saturated aqueous sodium bicarbonate (100ml). The aqueous phase was
15 extracted with further ethyl acetate (50ml) after pH adjustment (to pH ca 8). Combined extracts were washed with brine (100ml) dried (MgSO₄) and evaporated to a brown foam, which was chromatographed on silica (Merck 9385) eluting with 2% then 5% methanol in dichloromethane. The required fractions were combined and evaporated to give the title compound as a light
20 brown foam (672mg), mass spectrum (thermospray +ve) m/e 419 (MH⁺).

Intermediate 133-{tert-Butoxycarbonyl-[2-hydroxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-amino}-[2S,3S]-2-phenylpiperidine-1-carboxylic acid tert-butyl ester

25 A solution of [2-hydroxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-([2S,3S]-2-phenylpiperidine-3-yl)amine (418mg, 1mmol) in dichloromethane (5ml) was treated with di-t-butyl dicarbonate (240mg, 1.1mmol) and the solution was stirred for 19h at room temperature. Further reagent was then added (240mg) and after a further day the solution was evaporated to dryness. The resulting gum was chromatographed on silica (Merck 9385; 70g) eluting with 3:1 then 1:1
30 cyclohexane:ethyl acetate. The required fractions were combined and evaporated to provide the title compound as a light brown foam (487mg), $\nu(\text{CHBr}_3)$ 1663 and 1690cm⁻¹ (carbamate C=O).

Intermediate 14

3-{tert-Butoxycarbonyl-[2-(cyclopropylmethoxy)-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-amino}-[2S,3S]-2-phenylpiperidine-1-carboxylic acid tert-butyl ester

A solution of 3-{tert-butoxycarbonyl-[2-hydroxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-amino}-[2S,3S]-2-phenylpiperidine-1-carboxylic acid tert-butyl ester (402mg) in dimethylformamide (1ml) containing a suspension of finely ground potassium carbonate (89mg) was stirred at room temperature and treated with cyclopropylmethyl bromide (529mg). After 11 days stirring at room temperature the mixture was partitioned between ethyl acetate (50ml) and 2M hydrochloric acid (50ml). The organic phase was washed with brine (2x50ml), dried (MgSO₄) and evaporated to give a brown gum. This was chromatographed on silica (Merck 9385) eluting with 7:1 then 5:1 cyclohexane:ethyl acetate to give the title compound as a white foam (104mg), mass spectrum (electrospray +ve) m/e 673. (MH⁺).

Intermediate 15

[2-(Fluoromethoxy)-5-(5-trifluoromethyl-tetrazol-1-yl)]-benzaldehyde

A solution of [2-hydroxy-5-(5-trifluoromethyl)-tetrazol-1-yl]-benzaldehyde (200mg, 0.77mmole) in dimethylformamide (1ml) containing a suspension of potassium carbonate (107mg, 0.77mmol) was stirred in an ice-bath and treated with bromofluoromethane (few mls). Further bromofluoromethane (few mls) was added after 1.5h and the reaction mixture was allowed to reach room temperature. After 17h it was partitioned between ethyl acetate (50ml) and 2M-hydrochloric acid (50ml) and brine (2x50ml), dried (MgSO₄) and evaporated to the title compound as an off-white solid (211mg), ν (CHBr₃) 1698cm⁻¹ (C=O aldehyde).

Similarly prepared:-

Intermediate 16

[2-(2-Cyclopentoxo)-5-(5-trifluoromethyl-tetrazol-1-yl)]-benzaldehyde

From [2-hydroxy-5-(5-trifluoromethyl-tetrazol-1-yl)]-benzaldehyde (250mg, 0.97mmole) and bromocyclopentane (434mg, 2.91mmole) to give the title compound as a pale yellow solid (240mg), ν (KBr powder) 1687cm⁻¹ (C=O aldehyde).

Example 1(2-Cyclopentoxy-5-tetrazol-1-ylbenzyl)-([2S, 3S]-2-phenylpiperidin-3-yl) amine dihydrochloride

To a solution of [2S]-phenylpiperidin-[3S]-ylamine (1.0mmol) and 2-cyclopentoxy-5-tetrazol-1-ylbenzaldehyde (0.91mmol) in dichloromethane (4ml) and acetic acid (1.5mmol), triacetoxyborohydride (1.30mmol) was added. After 18h the solvent was reduced and the residue partitioned between ethyl acetate (25ml) and 2M sodium carbonate solution (25ml). The aqueous was re-extracted with ethyl acetate (2 x 25ml) and the combined organics dried (Na₂SO₄) and reduced to an oil. This was dissolved in hot ethanol (8ml) and conc.HCl added (177μl). After 4h the crystals were harvested and dried in vacuo at 50° overnight to give the title compound (309mg).

δ (D₂O) 1.50 - 1.80 (6H, m), 1.80 - 2.00 (2H, m), 2.10 - 2.20 (2H, m), 2.30 (1H, m) 2.55 (1H, m), 3.35 (1H, m), 3.75 (1H, m), 4.00 (1H, m), 4.15 (1H, d J=18Hz), 4.40 (1H, d J=18Hz), 4.95 (1H, d J=5Hz), 7.15 (3H, m), 7.40 (3H, m), 7.64 (1H, d J=4Hz), 7.80 (1H, dd J=11, 4 Hz), 9.58 (1H, s).

Assay: Found: C, 58.12%; H, 6.34%; N, 17.28%; Cl, 14.1%. C₂₄H₃₀N₆O. 2HCl requires C, 58.65%; H, 6.56%; N, 17.10%; Cl, 14.4%.

Similarly prepared:-

Example 2(2-Cyclopropylmethoxy-5-tetrazol-1-ylbenzyl)-([2S, 3S]-2-phenylpiperidin-3-yl) amine dihydrochloride

From 2-cyclopropylmethoxy-5-tetrazol-1-ylbenzaldehyde (0.59mmol) and [2S]-phenylpiperidin-[3S]-ylamine (0.65mmol) to give the title compound (134mg).

δ (D₂O) 0.30 (2H, m), 0.65 (2H, m), 1.20 (1H, m), 2.00 - 2.60 (4H, m), 3.35 (1H, m), 3.65 - 3.90 (3H, m), 4.02 (1H, m), 4.21 (1H, d J=18Hz), 4.45 (1H, d J=18Hz), 4.98 (1H, d J=5Hz), 7.15 (1H, d J=12Hz), 7.25 (2H, m), 7.45 (3H, m), 7.62 (1H, m), 7.82 (1H, dd J=11, 4Hz), 9.57 (1H, s).

Assay: Found: C, 57.70%; H, 6.32%; N, 17.43%; Cl, 14.7%. C₂₃H₂₈N₆O. 2HCl requires

C, 57.86%; H, 6.33%; N, 17.60%; Cl, 14.9%.

Example 3[2-(Cyclopropylmethoxy)-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-([2S,3S]-2-phenylpiperidin-3-yl)amine dihydrochloride

A solution of 3-{tert-butoxycarbonyl-[2-(cyclopropylmethoxy)-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-amino}-[2S,3S]-2-phenylpiperidine-1-carboxylic acid tert-butyl ester (101mg, 0.15mmole) in 4M-hydrogen chloride in dioxan (5ml) was kept at 44° for 3.25h and then evaporated. The residue was triturated with ethanol to give the title compound as a white solid (66mg), mass spectrum (thermospray +ve) m/e 473 (MH⁺). R_F 0.5 (100: 8:1 dichloromethane: ethanol: ammonia).

Example 4[2-(Fluoromethoxy)-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-([2S,3S]-2-phenylpiperidin-3-yl)amine dihydrochloride

A suspension of ([2S, 3S]-2-phenylpiperidin-3-yl)amine dihydrochloride (90mg, 0.36mmole) in dichloromethane (2ml) containing [2-(fluoromethoxy)-5-(5-trifluoromethyl-tetrazol-1-yl)]-benzaldehyde (104mg, 0.36mmole) was stirred at room temperature and treated with triethylamine (73mg, 0.72mmole) to give a solution. Glacial acetic acid (22mg, 0.36mmole) and sodium triacetoxyborohydride (114mg, 0.54mmole) were then added. After 2.75h the reaction mixture was evaporated to dryness. The residue was partitioned between ethyl acetate (50ml) and saturated aqueous sodium bicarbonate (50ml). The aqueous phase was further extracted with ethyl acetate (25ml). Combined extracts were washed with brine (2x50ml), dried (MgSO₄) and evaporated to a yellow gum. This was dissolved in hot ethanol (95% ca 3ml) and the solution was treated with concentrated hydrochloric acid (0.71mmole) to give white crystals of the title compound (107mg). Found C, 48.41; H, 4.91; N, 15.68; F, 14.52; C₂₁H₂₂F₄N₆O.2HCl (523.4) requires C, 48.19; H, 4.62; N, 16.02%; F, 14.26% Mass spectrum (thermospray +ve) m/e 451 (MH⁺).

Similarly prepared:-

Example 5[2-(2-Cyclopentoxo)-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-([2S,3S]-2-phenylpiperidin-3-yl)amine dihydrochloride.

From [2-(cyclopentoxy)-5-(5-trifluoromethyl-tetrazol-1-yl)]-benzaldehyde (100mg, 0.32mmole) and ([2S,3S]-2-phenylpiperidin-3-yl)amine (54mg, 0.32mmole) to give the title compound as a cream solid (114mg), mass spectrum (thermospray +ve) m/e 487 (MH⁺). R_F 0.41 (100:8:1 dichloromethane: ethanol: ammonia).

Example 6

a) 3-{tert-Butoxycarbonyl-[2-(2-methoxy-1-ethoxy)-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-amino}-[2S,3S]-2-phenylpiperidine-1-carboxylic acid tert-butyl ester

A solution of 3-{tert-butoxycarbonyl-[2-hydroxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-amino}-[2S,3S]-2-phenylpiperidine-1-carboxylic acid tert-butyl ester (177mg) in dimethylformamide (0.5ml) containing a suspension of finely ground potassium carbonate (39mg) was stirred and treated with 2-chloroethylmethyl ether (280mg) and potassium iodide (few crystals). The mixture was stirred at 44° for 9 days then at 60° for 2 days. The mixture was partitioned between ethyl acetate (25ml) and 2M hydrochloric acid (25ml). The aqueous phase was extracted with ethyl acetate (25ml) and the combined organic phases were washed with water (25ml) and brine (2x25ml), dried (MgSO₄) and evaporated to give a brown gum. This was chromatographed on silica (Merck 9385) eluting with 3:1 cyclohexane:ethyl acetate to give the title compound as a white foam (57mg), mass spectrum (thermospray + ve) m/e 677 MH⁺. The title compound was deprotected according to conventional procedures to give:-

b) [2-(2-methoxy-1-ethoxy)-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-([2S,3S]-2-phenylpiperidine-3-yl)amine

A solution of 3-{tert-Butoxycarbonyl-[2-(2-methoxy-1-ethoxy)-5-(trifluoromethyl-tetrazol-1-yl)-benzyl]-amino}-[2S,3S]-2-phenylpiperidine-1-carboxylic acid tert-butyl ester (54mg, 0.08mmol) in 4M hydrogen chloride in dioxan (5ml) was kept at 44° for 20 h and then evaporated to a white foam (48mg). Trituration of this material with ether followed by drying at 40° under vacuum gave the title compound (36mg)

δ (D₂O) 2.02 (2H, m), 2.18-2.38 and 2.38-2.58 (2H, 2m), 3.23-3.40 (1H, m), 3.43 (3H, s), 3.65-3.85 (3H, m), 3.91-4.25 (4H, m), 4.43 (1H, d, J=12Hz), 4.94 (1H, d, J=3Hz), 7.14 (1H, d, J=8Hz), 7.25 (2H, d, J=7Hz), 7.39-7.58 (4H, m), 7.69 (1H, dd, J=8, 2Hz).

Assay: Found: C, 49.15%; H, 5.25%; N, 14.40%. $C_{23}H_{27}F_3N_6O.C.7H_2O$ requires C, 49.15%; H, 5.45%; N, 14.95%.

Pharmacy Examples

Example A

5 STERILE FORMULATION

	mg/ml
Compound of the invention	0.3mg
Sodium Chloride USP	6.0mg
Sodium Acetate USP	2.6mg
Acetic Acid	1.1mg
Water for Injection USP qs to	1ml

The components are dissolved in a portion of the water for injection and the solution made up to final volume to provide 0.25mg/ml of the compound of the invention.

- 10 The solution may be packaged for injection, for example by filling and sealing in ampoules, vials or syringes. The ampoules, vials or syringes may be aseptically filled and/or terminally sterilised by, for example, autoclaving at 121°C.

TABLETS FOR ORAL ADMINISTRATION

- 15 Tablets may be prepared by the normal methods such as direct compression or wet granulation.

The tablets may be film coated with suitable film forming materials, such as Opadry White, type YS-1-7027, using standard techniques. Alternatively the tablets may be sugar coated.

Example B

Direct Compression Tablet	mg/Tablet
Compound of the invention	0.6mg
Magnesium Stearate	0.75mg
Avicel PH102 qs	150.00mg

The compound of the invention is passed through a 30 mesh sieve and blended with Avicel PH102 and magnesium stearate. The resultant mix is compressed into tablets using a suitable tablet machine fitted with 9/32" diameter punches.

Example C

WET GRANULATION

A formulation as described in Example B may be used. The compound of the invention (dihydrochloride) is dissolved in a suitable volume of granulating solution (purified water or 10% PVP K29/32 in water). After drying, the granules are screened, for example through 20 mesh screen, and blended with magnesium stearate. The granules are then compressed into tablets as described in Example B.

Example D

SUPPOSITORY

Compound of the invention	10.0mg
Witepsol W32, hard fat qs	2000.0mg

Blend micronized drug in a portion of the melted Witepsol W32 at approximately 36°C for approximately 15 minutes in a high speed mixer. Incorporate the homogenized slurry into the remaining portion of the melted Witepsol W32 and blend at approximately 36°C until satisfactory dispersion is achieved. Fill moulds with 2000mg formulation.

Example E

CAPSULE

	mg/capsule
Compound of the invention	12.0mg
Polyethylene glycol	92.89mg
Propylene glycol qs	200mg

Blend together polyethylene glycol and propylene glycol using heat as necessary. Stir until homogeneous. Add micronised compound of the invention to blend. Mix until homogenous. Fill into an appropriate gelatin mass to give soft gelatin capsules containing 200mg of the formulation.

Example FORAL SYRUP


	mg/ml
Compound of the invention	6.0mg
Sucrose	200mg
Methylparaben	1.2mg
Propylparaben	0.15mg
Flavouring	1.5mg
Citric Acid	0.1mg
Purified Water qs	1 ml

5

Dissolve the parabens in a small portion of the water that has been heated to approximately 90°C. Add the paraben solution to a large portion of the remaining water with mixing. Add and dissolve the other components. Bring the formulation to final volume and mix until homogenous. Fill the formulation into a container, such as a unit dose cup or a bottle for multiple-dose use.


Claims

1. A compound of formula (I)



wherein R^1 is $-O-(CH_2)_pC_{3-7}$ cycloalkyl, $-O-C_{1-7}$ fluoroalkyl, or $-O-(CH_2)_nX$;

5 R^2 is



R^3 is a hydrogen or halogen atom;

R^4 and R^5 may each independently represent a hydrogen or halogen atom, or a C_{1-4} alkyl, C_{1-4} alkoxy or trifluoromethyl group;

10 X is selected from $C(O)-NR^7R^8$, $C(O)R^9$, NR^7R^8 , $SO_2NR^7R^8$, $NHSO_2R^9$, $S(O)_2R^9$, OC_{1-4} alkyl, NO_2 , CO_2H , CO_2C_{1-4} alkyl, CN or, when n is 2, X may also represent OH , SH or NH_2 ;

R^6 is a hydrogen atom, a C_{1-4} alkyl, $(CH_2)_m$ cyclopropyl, $-S(O)_2C_{1-4}$ alkyl, phenyl, $NR^{10}R^{11}$, $CH_2C(O)CF_3$, trifluoromethyl, difluoromethyl or cyano group;

15 R^7 and R^8 may each independently represent hydrogen atoms or a C_{1-4} alkyl group;

R^9 represents a C_{1-4} alkyl or trifluoromethyl group;

R^{10} and R^{11} may each independently represent a hydrogen atom, or a C_{1-4} alkyl or acyl group;

x represents zero or 1;

n is 1 or 2;

s represents zero, 1 or 2;


m represents zero or 1;

5 p represents zero or 1;

or a pharmaceutically acceptable salt or solvate thereof.

2. A compound according to Claim 1 wherein

R² is



10 x is zero and R³, R⁴ and R⁵ are each hydrogen atoms.

3. A compound according to Claim 1 or Claim 2 wherein R¹ is a cyclopropylmethoxy, cyclopentyloxy, difluoromethyloxy, trifluoromethyloxy, 2,2,2-trifluoroethyloxy, fluoromethyloxy, OCH₂OMe or O-CH₂CH₂OMe group.

15 4. A compound according to any one of Claims 1 to 3 wherein R⁶ is a hydrogen atom or a C₁₋₄alkyl or a trifluoromethyl group.

5. A compound selected from

(2-Cyclopentoxy-5-tetrazol-1-ylbenzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

(2-Cyclopropylmethoxy-5-tetrazol-1-ylbenzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

20 (2-Cyclopentoxy-5-(5-trifluoromethyltetrazol-1-yl)benzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

(2-Cyclopropylmethoxy-5-(5-trifluoromethyltetrazol-1-yl)benzyl)-([2S,3S]-2-

phenylpiperidin-3-yl)amine;

(2-Fluoromethoxy-5-(5-trifluoromethyltetrazol-1-yl)benzyl)-([2S,3S]-2-phenylpiperidin-3-yl)amine;

[2-(2-methoxy-1-ethoxy)-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-([2S,3S]-2-phenylpiperidine-3-yl)amine

or a pharmaceutically acceptable salt or solvate thereof.

6. A compound according to any one of Claims 1 to 5 for use in therapy.


7. A pharmaceutical composition comprising a compound according to any one of Claims 1 to 5 or a pharmaceutically acceptable salt or solvate thereof, together with a pharmaceutically acceptable carrier.

8. A method for the treatment of a condition mediated by tachykinins, including substance P and other neurokinins, in a mammal comprising administration of an effective amount of a compound according to any one of claims 1 to 5 or a pharmaceutically acceptable salt or solvate thereof.

9. The use of a compound according to any one of claims 1 to 5, or a pharmaceutically acceptable salt or solvate thereof, in the preparation of a medicament for use in the treatment of conditions mediated by tachykinins, including substance P and other neurokinins.


10. A process for preparing a compound of formula (I) as defined in Claim 1, or a pharmaceutically acceptable salt or solvate thereof which comprises:

(A) reacting a compound of formula (II)



with a compound of formula (III)


44



followed by reduction: or


(B) a process for preparing a compound of formula (I) where R^2 is a group (A) as defined hereinbefore, x is zero and R^6 is $-NH_2$ which comprises reacting a

5 compound of formula (XXIV)




with ammonium chloride and sodium azide: or

(C) reducing a compound of formula (XXV)



10 (D) a process for preparing compounds of formula (I), excluding compounds where R^1 represents cyclopropyloxy, cyclobutyloxy and trifluoromethyloxy which comprises reacting a compound of formula (XXVII)


45



or an amino-protected derivative thereof, with $L-(CH_2)_pC_{3-7}$ cycloalkyl or $L-C_{1-7}$ fluoroalkyl where p is as defined hereinbefore and L represents a suitable leaving group in the presence of a base followed by deprotection where necessary: or

5

(E) a process for preparing compounds of formula (I) where R^2 is a group A as defined above and R^6 represents cyano which comprises reacting a compound of formula (XXIX)



10

where L represents a suitable leaving group, with potassium cyanide.

INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/EP 96/00082A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D401/12 A61K31/445

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	J. MED. CHEM. (JMCMAR,00222623);95; VOL.38 (26); PP.4985-92, MEDICINES RESEARCH CENTRE;DEPARTMENT OF MEDICINAL CHEMISTRY; STEVENAGE/ HERTS; UK (GB), WARD P ET AL 'Discovery of an Orally Bioavailable NK1 Receptor Antagonist, (2S,3S)-(2-Methoxy-5-tetrazol-1-ylbenzyl)(2-phenylpiperidin-3-yl)amine (GR203040), with Potent Antiemetic Activity' see table 2, page 4988 ---	1-10
P,X	WO,A,95 08549 (GLAXO GROUP LTD.;UK) 30 March 1995 see the whole document ---	1-10
A	WO,A,93 01170 (PFIZER INC.;USA) 21 January 1993 see page 5 - page 6 ---	1-10
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- 'A' document defining the general state of the art which is not considered to be of particular relevance
- 'E' earlier document but published on or after the international filing date
- 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- 'O' document referring to an oral disclosure, use, exhibition or other means
- 'P' document published prior to the international filing date but later than the priority date claimed

- 'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- 'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- 'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- '&' document member of the same patent family

Date of the actual completion of the international search

16 April 1996

Date of mailing of the international search report

10. 05. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Steendijk, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 96/00082

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO,A,93 00331 (PFIZER INC.;USA) 7 January 1993 see page 9 - page 11 -----</p>	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Appl. No.

PCT/EP 96/00082

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9508549	30-03-95	AU-B- 7697494	10-04-95
WO-A-9301170	21-01-93	US-A- 5232929	03-08-93
		CA-A- 2111461	21-01-93
		EP-A- 0594636	04-05-94
		JP-T- 6508828	06-10-94
		US-A- 5332817	26-07-94
WO-A-9300331	07-01-93	AU-B- 657967	30-03-95
		AU-B- 1889392	25-01-93
		BR-A- 9206161	31-10-95
		CA-A- 2109613	07-01-93
		CN-A- 1067655	06-01-93
		CZ-A- 9203908	13-04-94
		DE-U- 9290083	17-02-94
		EP-A- 0589924	06-04-94
		HU-A- 67434	28-04-95
		HU-A- 70499	30-10-95
		JP-B- 7110850	29-11-95
		JP-T- 6506473	21-07-94
		NO-A- 934691	17-12-93
		SK-A- 390892	13-09-95
		ZA-A- 9204528	20-12-93